

FORM PTO-1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK
 OFFICE (MODIFIED)

ATTORNEY'S DOCKET NUMBER

X-11869

**TRANSMITTAL LETTER TO THE UNITED STATES
 DESIGNATED/ELECTED OFFICE (DO/EO/US)
 CONCERNING A FILING UNDER 35 U.S.C. 371**

U.S. APPLICATION NO. (if known, see 37 C.F.R. 1.5)

09/868974 ✓

INTERNATIONAL APPLICATION NO

PCT/US99/30395 ✓

INTERNATIONAL FILING
 DATE

December 21, 1999 ✓
 (12.21.1999)

PRIORITY DATE CLAIMED

December 22, 1998 (12.22.1998) ✓

TITLE OF INVENTION: SHELF-STABLE FORMULATION OF GLUCAGON-LIKE PEPTIDE-1

APPLICANT(S) FOR DO/EO/US: Mark Laurence Brader, et al.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☐ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☒ A copy of the International Preliminary Examination Report (IPER), including any annexes, and, if not in English, an English language translation of the annexes to the IPER under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☐ Other items or information:

[PAGE 2 OF 2]

09868974-062001

JCO5 Rec'd PCT/PTO 26 SEP 2001

Saco
CERTIFICATE OF MAILING

09/868974

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231, on the date appearing below.
ELI LILLY AND COMPANY

By *Hinda M. Dunbar*

Date *September 18, 2001*

PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Mark L. Brader, et al.)
Serial No. : 09/868,974)
Filed : December 21, 1999)
For : SHELF-STABLE FORMULATION OF)
GLUCAGON-LIKE PEPTIDE-1)
Docket No. : X-11869)

STATEMENT TO SUPPORT FILING AND SUBMISSION IN ACCORDANCE

WITH 37 C.F.R. 1.821(f) (SEQUENCE LISTING)

Assistant Commissioner for Patents

Washington, D. C. 20231

Sir:

I hereby affirm that the content of the paper and computer readable copies of the Sequence Listing, submitted in accordance with 37 C.F.R. 1.821(c) and (e), respectively, are the same and include no new matter.

AMENDMENT DIRECTING ENTRY OF SEQUENCE LISTING

INTO THE SPECIFICATION

Please amend the application by adding the Sequence Listing submitted herewith.

Serial No. 09/868,974

-2-

Respectfully submitted,



Mark J. Stewart, Ph.D.
Attorney for Applicants
Registration No. 43,936
Phone: 317-276-0280

Eli Lilly and Company
Patent Division/MJS
Lilly Corporate Center
Indianapolis, Indiana 46285

September 18, 2001

SHELF-STABLE FORMULATION OF GLUCAGON-LIKE PEPTIDE-1

Background of the Invention

Glucagon-like peptide-1 (7-37)-OH (GLP-1) is a 31 amino acid hormone that is produced by post-translational processing of the preglucagon gene product in the brain, stomach, intestine, and pancreas. The main physiological function of GLP-1 is to regulate insulin secretion in response to glucose, and thus it has the ability to normalize blood glucose levels. As such, there has been interest in GLP-1, its analogs and derivatives as potential therapeutic agents for the treatment of diabetes. A particular advantage to the use of GLP-1 over other drugs in the treatment of diabetes is that administration of GLP-1 at doses in the 1-5 nmole range exhibit few adverse side effects, such as hypoglycemia. Unexpectedly, GLP-1 also has been shown to work in patients that have secondary failure to sulfonylurea drugs, the most common drug type for the treatment of type II diabetes. GLP-1 also is a potent inhibitor of gastric acid secretion and gastric emptying.

In general, effective therapeutic administration of peptides can be problematic since peptides often are degraded in the gastrointestinal tract by various peptidases. Additionally, certain peptide treatment protocols require either continuous or repeated administration of the peptide agent over an extended period of time. Repeated injections cause both inconvenience and discomfort to the user. Thus, chronic use of the peptide agent, which would be required for patients afflicted with diabetes, would result in inconvenience and discomfort to the user.

The long-term stability of peptides, particularly GLP-1, as components of a pharmaceutical composition for administration to mammals, is questionable. Such a lack of stability adversely affects bioavailability. In fact, when stored at low temperatures of 4° C, by-products of GLP-1(7-37) have been found as early as eleven months after sample preparation (see Mojssov, *Int. J. Peptide Protein Res.*, Vol. 40, pages 333-343 (1992)). Additionally, the biological half-life of GLP-1 molecules, particularly those molecules affected by the activity of dipeptidyl-peptidase IV (DPPIV), is quite short. For example,

WO 00/37098

PCT/US99/30395

the biological half-life of GLP-1(7-37) is only 3 to 5 minutes (see U.S. Patent No. 5,118,666), which is further augmented by its rapid absorption following parenteral administration to a mammal.

Another factor decreasing the bioavailability of GLP-1 is the solubility of GLP-1 when incorporated into an aqueous solution. The solubility of GLP-1 is highly dependent on the environment, such as the choice of buffering system, and the treatment that the peptide has undergone. For example, conversion of a peptide into its salt form plays a role in its solubility. In this regard, synthetic GLP-1 is highly soluble in neutral phosphate buffered saline. Because the solubility of the peptide is high in such aqueous solutions, slow release of the peptide can be difficult to attain unless the peptide is incorporated into a system for slow release.

Stable formulations of therapeutic agents are particularly required for use in delivery devices that expose these agents to elevated temperatures and/or mechanical stress. For example, stable GLP-1 formulations are required for use in continuous infusion systems and pen delivery devices. Current formulations provide only limited stability in these types of delivery devices.

In continuous infusion systems, a fluid containing a therapeutic agent is pumped from a reservoir, usually to a subcutaneous, intravenous, or intraperitoneal depot. The reservoir, which must be refilled periodically, is attached to the patient's body, or is implanted in the patient's body. In either case, the patient's body heat and body motion, and turbulence in the tubing and pump impart a relatively high amount of thermo-mechanical energy to the formulation. In the interest of minimizing the frequency with which the reservoir is refilled, and of minimizing the size of the reservoir, formulations having a relatively high concentration of the therapeutic agent are advantageous.

Injector pens also have been developed to allow diabetic patients to accurately measure and administer controlled doses of insulinotropic agents. Generally, these pens are secured onto a cartridge having a particular quantity of liquid medication sealed therein. The cartridge includes a plunger and a mechanism for advancing the plunger in the cartridge in such a manner to dispense the medication. Injector pens may be reusable or disposable. In reusable pens, a user can change a spent cartridge and reset the leadscrew of the pen back to its initial position. In a disposable pen, the cartridge is permanently

WO 00/37098

PCT/US99/30395

embodiment, the GLP-1 molecule of the formulation is a derivative of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:

$\text{NH}_2\text{-His}^7\text{-Ala-Glu-Gly}^{10}\text{-Thr-Phe-Thr-Ser-Asp}^{15}\text{-Val-Ser-Ser-Tyr-Leu}^{20}\text{-Glu-Gly-Gln-Ala-Ala}^{25}\text{-Lys-Glu-Phe-Ile-Ala}^{30}\text{-Trp-Leu-Val-X}$ (SEQ ID NO:3)

and a pharmaceutically-acceptable salt thereof, wherein X is selected from the group consisting of Lys and Lys-Gly; a pharmaceutically-acceptable lower alkylester of the peptide; and a pharmaceutically-acceptable amide of the peptide selected from the group consisting of amide, lower alkyl amide, and lower dialkyl amide. In another preferred embodiment, the formulation also comprises a long-acting insulin agent.

The present invention also provides a method of enhancing the expression of insulin in a mammalian pancreatic β -type islet cell in need of such enhancement, comprising administering to the cell, an effective amount of a shelf-stable pharmaceutical formulation, wherein the formulation comprises a therapeutically effective amount of a GLP-1 molecule, a pharmaceutically acceptable preservative, and a tonicity modifier, and wherein the formulation has a pH that is about 8.2 to about 8.8. In a preferred embodiment, the formulation used in the therapeutic method comprises a buffer, such as TRIS. In another preferred embodiment, the formulation used in the therapeutic method further comprises a surfactant, such as Brij-35. In an additional preferred embodiment, the GLP-1 molecule of the formulation thus administered is an analog of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:

$\text{R}_1\text{-X-Glu-Gly}^{10}\text{-Thr-Phe-Thr-Ser-Asp}^{15}\text{-Val-Ser-Ser-Tyr-Leu}^{20}\text{-Y-Gly-Gln-Ala-Ala}^{25}\text{-Lys-Z-Phe-Ile-Ala}^{30}\text{-Trp-Leu-Val-Lys-Gly}^{35}\text{-Arg-R}_2$ (SEQ ID NO:2)

and a pharmaceutically-acceptable salt thereof, wherein R_1 is His or desamino-histidine, X is Ala, Gly or Val, Y is Glu or Gln, Z is Glu or Gln and R_2 is Gly-OH. In an especially preferred embodiment, the GLP-1 molecule administered is according to SEQ ID NO: 2, wherein R_1 is L-histidine, X is Val, Y is Glu, Z is Glu, and R_2 is Gly-OH. In an alternative preferred embodiment, the GLP-1 molecule administered is a derivative of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:

$\text{NH}_2\text{-His}^7\text{-Ala-Glu-Gly}^{10}\text{-Thr-Phe-Thr-Ser-Asp}^{15}\text{-Val-Ser-Ser-Tyr-Leu}^{20}\text{-Glu-Gly-Gln-Ala-Ala}^{25}\text{-Lys-Glu-Phe-Ile-Ala}^{30}\text{-Trp-Leu-Val-X}$ (SEQ ID NO:3)


and a pharmaceutically-acceptable salt thereof, wherein X is selected from the group consisting of Lys and Lys-Gly; a pharmaceutically-acceptable lower alkylester of the peptide;

WO 00/37098


PCT/US99/30395

and a pharmaceutically-acceptable amide of the peptide selected from the group consisting of amide, lower alkyl amide, and lower dialkyl amide.

The present invention also provides a method of treating diabetes comprising administering to a patient in need of such treatment an effective amount of a shelf-stable pharmaceutical formulation, wherein the formulation comprises a therapeutically effective amount of a GLP-1 molecule, a pharmaceutically acceptable preservative, and a tonicity modifier, and wherein the formulation has a pH that is about 8.2 to about 8.8. In a preferred embodiment, the formulation used in the therapeutic method comprises a buffer, such as TRIS. In another preferred embodiment, the formulation used in the therapeutic method further comprises a surfactant, such as Brij-35. In an additional preferred embodiment, the GLP-1 molecule of the formulation thus administered is an analog of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:



and a pharmaceutically-acceptable salt thereof, wherein R_1 is His or desamino-histidine, X is Ala, Gly or Val, Y is Glu or Gln, Z is Glu or Gln and R_2 is Gly-OH. In an especially preferred embodiment, the GLP-1 molecule administered is according to SEQ ID NO: 2, wherein R_1 is L-histidine, X is Val, Y is Glu, Z is Glu, and R_2 is Gly-OH. In an alternative preferred embodiment, the GLP-1 molecule administered is a derivative of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:



and a pharmaceutically-acceptable salt thereof, wherein X is selected from the group consisting of Lys and Lys-Gly; a pharmaceutically-acceptable lower alkylester of the peptide; and a pharmaceutically-acceptable amide of the peptide selected from the group consisting of amide, lower alkyl amide, and lower dialkyl amide.

An additional embodiment encompasses a method of providing meal-time glycemic control and basal glycemic control with a single injection comprising administering to a patient in need thereof an effective amount of a shelf-stable pharmaceutical formulation, wherein the formulation comprises a therapeutically effective amount of a GLP-1 molecule, a long acting insulin agent, a pharmaceutically acceptable preservative, and a tonicity modifier, wherein said formulation has a pH that is about 8.2 to about 8.8.

WO 00/37098

PCT/US99/30395


captured in the pen which is disposed of after the contents of the cartridge have been exhausted.

With the development of GLP-1 as well as analogs and derivatives thereof for the treatment of diabetes, there exists a need to improve treatment regimes that can balance chemical and physical stability with chronic use requirements of diabetic patients.

Summary of the Invention

In order to overcome the problems of chemical and physical stability of GLP-1 formulations, the present inventors have developed a shelf-stable formulation of GLP-1. In particular, the inventors have discovered that when certain physiologically tolerated buffers are used in formulations of GLP-1, the physical stability of such formulations is unexpectedly and considerably greater than when compared to GLP-1 formulations made with a phosphate buffer. In addition, maintaining the pH in a range of about 8.2 to about 8.8 unexpectedly improves the chemical stability of the formulations. The shelf-stable formulation of GLP-1 of the invention comprises a therapeutically effective amount of a GLP-1 molecule, a pharmaceutically acceptable preservative, and a tonicity modifier, wherein the pH of said formulation is maintained in the range from about 8.2 to about 8.8.

In accordance with the chemical and physical stability needs of GLP-1 formulations, the present invention provides a shelf-stable pharmaceutical formulation comprising a therapeutically effective amount of a GLP-1 molecule, a pharmaceutically acceptable preservative, and a tonicity modifier, wherein the formulation has a pH that is about 8.2 to about 8.8. In a preferred embodiment, the formulation further includes a buffer, such as TRIS. In another preferred embodiment, the formulation comprises a surfactant, such as Brij-35. In an additional preferred embodiment, the GLP-1 molecule of the formulation is an analog of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:



and a pharmaceutically-acceptable salt thereof, wherein R_1 is His or desamino-histidine, X is Ala, Gly or Val, Y is Glu or Gln, Z is Glu or Gln and R_2 is Gly-OH. In an especially preferred embodiment, the GLP-1 molecule is according to SEQ ID NO: 2, wherein R_1 is L-histidine, X is Val, Y is Glu, Z is Glu, and R_2 is Gly-OH. In an alternative preferred

Detailed Description of the Preferred Embodiments

The present invention provides a shelf-stable formulation of GLP-1, GLP-1 analogs, and GLP-1 derivatives. Because it is known that there are problems with long term stability of GLP-1 as a component in a pharmaceutical composition, the present inventors developed a pharmaceutical formulation which stabilizes GLP-1, its derivatives and analogs. This development led to the shelf-stable GLP-1 formulations of the invention.

In another embodiment, the present invention encompasses a GLP-1 formulation that also comprises a long acting diabetic agent. It has long been a goal of insulin therapy to mimic the pattern of endogenous insulin secretion in normal individuals. The daily physiological demand for insulin fluctuates and can be separated into two phases: (a) the absorptive phase requiring a pulse of insulin to dispose of the meal-related blood glucose surge, and (b) the post-absorptive phase requiring a sustained delivery of insulin to regulate hepatic glucose output for maintaining optimal fasting blood glucose. Accordingly, effective therapy for people with diabetes generally involves the combined use of two types of exogenous insulin formulations: a fast-acting meal time insulin provided by bolus injections and a long-acting, so-called, basal insulin, administered by injection once or twice daily to control blood glucose levels between meals. Examples of commercial basal insulin preparations include NPH (Neutral Protamine Hagedorn) insulin, protamine zinc insulin (PZI), and ultralente (UL).

The term "stability" is used to mean chemical as well as physical stability. Physical stability refers to properties such as protein aggregation, which can be measured by a sample's attenuation of light. The measurement relates to the turbidity of a formulation. Turbidity is produced by aggregation or precipitation of proteins or complexes in the formulation and is indicative of decreased stability of a solution formulation. The more turbid a protein preparation, the less stable the preparation is. Stability also refers to the chemical stability of the formulation such as the propensity of the proteins to form high order polymers which is indicative of decreased stability.

One factor that plays a role in the stability of GLP-1 formulations is the maintenance of pH at a prescribed level. Specifically, the present inventors have found that achieving and maintaining the pH of the formulation at about 8.2 to about 8.8 is advantageous. Typical peptide formulations have a more neutral pH of 7 to about 7.8 or an acidic pH. Furthermore, a composition containing a GLP-1 molecule that has a pH in the range of

about 6.8 to about 7.5 exhibits less physical stability than a composition of a GLP-1 molecule containing a preservative and having a pH in the range of about 8.2 to about 8.8. A preserved formulation which has a pH of less than about 8.0 tends to exhibit turbidity, a telltale sign of decreased physical stability of the peptide formulation. Conversely, a formulation which has a pH greater than about 8.8 tends to have decreased chemical stability.

Therefore, the invention contemplates GLP-1 formulations having a pH range of about 8.2 to about 8.8, which preserves optimal chemical and physical stability of the GLP-1 molecule. A particularly preferred range for the inventive GLP-1 formulations is about 8.3 to about 8.6, and a most particularly preferred pH range is about 8.4 to about 8.5. As used in this specification with respect to pH, the term "about" means plus or minus 0.1 pH units. Thus, a pH of "about 8.5" denotes a pH of 8.4 to 8.6.

GLP-1 molecules themselves exhibit a buffering capacity. However, to maintain the pH of the composition for long term storage and stability, it is preferable to add a buffer.

The choice of buffer affects the chemical and physical stability of the formulation because it influences pH. There are very few pharmaceutically acceptable buffers in the alkaline range. Phosphate buffers, which are typically used in peptide pharmaceutical formulations, cannot maintain a pH range of 8.2 to about 8.8. However, the present inventors have discovered that certain other amine-containing buffers are capable of imparting chemical as well as physical stability to formulations of GLP-1 molecules.

The buffers used in the present invention preferably provide a buffering capacity in the range of about 8.2 to about 8.8. The buffers which are used may be tromethane (TRIS), and amino-acid based buffers such as lysine and hydroxy-lysine. Although any non-phosphate buffer which has a buffering capacity in the range of about 8.2 to about 8.8 may be used, TRIS is the preferred buffer for the formulations of the present invention. The term "TRIS" refers to 2-amino-2-hydroxymethyl-1,3-propanediol (also known in the art as tromethane, trimethylol aminomethane or tris(hydroxymethyl) aminomethane), and to any pharmacologically acceptable salt thereof. The free base and the hydrochloride form are two common forms of TRIS. TRIS is one of the few buffers which is capable of maintaining the pH at the alkaline level desired, thereby stabilizing the formulation.

A second factor that plays a role in the stability of GLP-1 formulations is the concentration of the GLP-1 molecule that is used in the inventive formulation. The present

WO 00/37098

PCT/US99/30395

inventors determined that a concentration of about 0.30 to about 0.65 mg/ml of the GLP-1 molecule was stable in the inventive formulations. However, a concentration which was about equal to or greater than 1 mg/ml was unstable. This stability was evidenced by the development of turbidity in the formulation. A particularly stable formulation includes about 0.5 mg/ml of a GLP-1 molecule.

An additional factor which contributes to the overall stability of the GLP-1 formulations of the present invention is the choice of preservative. The preservative is an essential component in the formulation because it enables multiple uses of the formulation. While it is typical that most preservatives would be capable of stabilizing a pharmaceutical formulation, some pharmaceutically acceptable preservatives act to promote physical instability of the formulation. The present inventors have found that a phenolic preservative is preferred. Specifically, the preservative may be m-cresol, phenol, benzyl alcohol, or methyl paraben and is present in an amount from about 2 mg/ml to about 6 mg/ml. Ultimately, the concentration of preservative necessary for effective preservation depends on the preservative used, the pH of the formulation, and whether substances that bind or sequester the preservative are also present. Preferably, m-cresol is used in the formulations as a preservative.

While a buffer and a preservative are most preferably included in the formulation, other additional excipients may be included, such as a tonicity modifier and/or a surfactant as well as distilled water for injections.

The tonicity modifier may be included to make the formulation approximately isotonic with bodily fluid depending on the mode of administration. The concentration of the tonicity modifier is in accordance with the known concentration of a tonicity modifier in a peptide formulation. A preferable tonicity modifier used in the present invention is glycerol.

A surfactant, which may be included in the formulation of the present invention, can be cationic, anionic, or non-ionic. A preferable class of surfactants is polyoxyethylene ethers. A preferred surfactant useful in the present invention is Brij-35, a polyoxyethylene 23 lauryl ether, available from ICI United States, Inc.

The present invention contemplates use of not only natural GLP-1, but also analogs, derivatives and salts of GLP-1. As used herein, the term "a GLP-1 molecule" refers (1) to the naturally-occurring GLP-1, which is GLP-1 (7-37)-OH; (2) GLP-1(7-36)NH₂, as well


WO 00/37098

PCT/US99/30395

as (3) GLP-1 (7-37); (4) natural and synthetic functional GLP-1 analogs; (5) derivatives of GLP-1 and (6) salts of any of the aforementioned molecules.

A "GLP-1 analog" is defined as a molecule having one or more amino acid substitutions, deletions, inversions, or additions relative to GLP-1(7-37) and may include the D-amino acid forms. Numerous GLP-1 analogs are known in the art and include, but are not limited to, GLP-1(7-34), GLP-1(7-35), GLP-1(7-36)NH₂, Gln⁹-GLP-1(7-37), d-Gln⁹-GLP-1(7-37), Thr¹⁶-Lys¹⁸-GLP-1(7-37), and Lys¹⁸-GLP-1(7-37), Gly⁸-GLP-1(7-36)NH₂, Gly⁸-GLP-1(7-37)OH, Val⁸-GLP-1(7-37)OH, Met⁸-GLP-1(7-37)OH, acetyl-Lys⁹-GLP-1(7-37), Thr⁹-GLP-1(7-37), D-Thr⁹-GLP-1(7-37), Asn⁹-GLP-1(7-37), D-Asn⁹-GLP-1(7-37), Ser²²-Arg²³-Arg²⁴-Gln²⁶-GLP-1(7-37), Arg²³-GLP-1(7-37), Arg²⁴-GLP-1(7-37), α-methyl-Ala⁸-GLP-1(7-36)NH₂, and Gly⁸-Gln²¹-GLP-1(7-37)OH, and the like.

Other GLP-1 analogs consistent with the present invention are described by the formula:



wherein: R₁ is selected from the group consisting of L-histidine, D-histidine, desamino-histidine, 2-amino-histidine, beta-hydroxy-histidine, homohistidine, alpha-fluoromethyl-histidine, and alpha-methyl-histidine; X is selected from the group consisting of Ala, Gly, Val, Thr, Ile, and alpha-methyl-Ala; Y is selected from the group consisting of Glu, Gln, Ala, Thr, Ser, and Gly; Z is selected from the group consisting of Glu, Gln, Ala, Thr, Ser, and Gly; and R₂ is selected from the group consisting of NH₂, and Gly-OH.

GLP-1 analogs also have been described in WO 91/11457, and include GLP-1(7-34), GLP-1(7-35), GLP-1(7-36), or GLP-1(7-37), or the amide form thereof, and pharmaceutically-acceptable salts thereof, having at least one modification selected from the group consisting of:

(a) substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, phenylalanine, arginine, or D-lysine for lysine at position 26 and/or position 34; or substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, phenylalanine, lysine, or a D-arginine for arginine at position 36;

(b) substitution of an oxidation-resistant amino acid for tryptophan at position 31;

WO 00/37098

PCT/US99/30395

(c) substitution of at least one of: tyrosine for valine at position 16; lysine for serine at position 18; aspartic acid for glutamic acid at position 21; serine for glycine at position 22; arginine for glutamine at position 23; arginine for alanine at position 24; and glutamine for lysine at position 26; and

(d) substitution of at least one of: glycine, serine, or cysteine for alanine at position 8; aspartic acid, glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine for glutamic acid at position 9; serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine for glycine at position 10; and glutamic acid for aspartic acid at position 15; and

(e) substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine, or the D- or N-acylated or alkylated form of histidine for histidine at position 7; wherein, in the substitutions described in (a), (b), (d), and (e), the substituted amino acids can optionally be in the D-form and the amino acids substituted at position 7 can optionally be in the N-acylated or N-alkylated form.

Preferred GLP-1 molecules used in the present inventive formulation also include analogs of GLP-1 (7-37)NH₂ and GLP-1 (7-37) in which one or more amino acids which are not present in the original sequence are added or deleted, and derivatives thereof. Specifically, His and desamino-histidine are preferred for R₁. Ala, Gly and Val are preferred at the "X" position. Also, Glu and Gln are preferred for at the "Y" position. Glu and Gln are preferred at the "Z" position and Gly-OH is preferred for R₂.

A particularly preferred GLP-1 analog is known as Val(8)GLP-1 (V8GLP-1) and has a formula according to SEQ ID NO:2, wherein R₁ is L-histidine, X is Val, Y is Glu, Z is Glu and R₂ is Gly-OH.

A "GLP-1 derivative" is defined as a molecule having the amino acid sequence of GLP-1(7-37) or of a GLP-1 analog, but additionally comprises chemical modification of one or more of its amino acid side groups, α -carbon atoms, terminal amino group, or terminal carboxylic acid group. A chemical modification includes, but is not limited to, adding chemical moieties, creating new bonds, and removing chemical moieties. Modifications at amino acid side groups include, without limitation, acylation of lysine ϵ -amino groups, N-alkylation of arginine, histidine, or lysine, alkylation of glutamic or aspartic carboxylic acid

WO 00/37098

PCT/US99/30395

"Biosynthetic GLP-1 analogs" are defined as any GLP-1 analogs or derivatives which contain only naturally occurring amino acid residues and are thus capable of being expressed by living cells, including recombinant cells and organisms.

Methods for preparing the GLP-1 molecules which are incorporated into the shelf stable formulations of the invention are well known to the skilled artisan. In one method, GLP-1 molecules are prepared by well-known methods of peptide synthesis, such as those described in Merrifield, (*Chem. Soc. Vol. 85*, page 2149, 1962). It is also contemplated that the molecules can be prepared by fragmenting the natural amino acid sequence of GLP-1(7-37) with, for example, a proteolytic enzyme. It is also contemplated that recombinant DNA techniques can be used to prepare the molecules, such as the methods of Maniatis *et al.* (*Molecular Biology: A Laboratory Manual*, CSH, 1982).

Administration may be via any route known to be effective by the physician of ordinary skill. Parenteral administration is preferred. Parenteral administration is commonly understood as administration by other than a gastro-intestinal route. Preferred parenteral routes for administering the formulations of the present invention include intravenous, intramuscular, subcutaneous, intraperitoneal, intraarterial, nasal, pulmonary, and buccal routes. Intravenous, intraperitoneal, intramuscular, and subcutaneous routes of administration of the compounds used in the present invention are more preferred parenteral routes of administration. Intravenous, intraperitoneal, and subcutaneous routes of administration of the formulations of the present invention yet more highly preferred. The most preferred route of administration is via a pen injection system, wherein either a 1.5 ml cartridge or a 3.0 ml cartridge is utilized.

Administration via certain parenteral routes may involve introducing the formulations of the present invention into the body of a patient through a needle or a catheter, propelled by a sterile syringe or some other mechanical device such as an continuous infusion system. A formulation provided by the present invention may be administered using a syringe, injector, pump, or any other device recognized in the art for parenteral administration. A formulation of the present invention may also be administered as an aerosol for absorption in the lung or nasal cavity. The formulations may also be administered for absorption through the mucus membranes, such as in buccal administration.

The amount of a formulation of the present invention that is administered to treat diabetes or hyperglycemia depends on a number of factors, among which are included, without limitation, the patient's sex, weight and age, the underlying causes of the condition or disease to be treated, the route of administration and bioavailability, the persistence of the administered GLP-1, its analog, or its derivative in the body, the formulation, and the potency of the GLP-1 molecule. Where administration is intermittent, the amount per administration should also take into account the interval between doses, and the bioavailability of the GLP-1 molecule from the formulation. Administration of the formulation of the present invention could be continuous. It is within the skill of the ordinary physician to titrate the dose and infusion rate or frequency of administration of the formulation of the present invention to achieve the desired clinical result.

The formulations of the present invention have insulintropic activity. Thus, another aspect of the present invention provides a method of enhancing the release of insulin from pancreatic islet cells comprising providing to a mammalian pancreatic β -type islet cell the inventive formulation which contains an effective amount of GLP-1, a GLP-1 analog, a GLP-1 derivative or a salt thereof.

That the present invention provides formulations of GLP-1, its analogs, and its derivatives having greatly increased physical and chemical stability relative to peptide formulations the art will be readily apparent from the following data and examples.

The following examples and preparations are provided merely to further illustrate the preparation of the formulations of the invention. The scope of the invention is not limited to the following examples.

Example 1-Preparation and chemical stability testing

66.2 mg of Val(8) GLP-1 was dissolved in water at 1.0 mg/ml and adjusted to pH 8.51. Three formulations were made as follows:

- (A) A 21.5 ml aliquot of peptide solution in water was mixed with 21.5 ml of 0.63% m-cresol-3.2% glycerol and the final pH was set to 8.48. The solution was passed thru a 0.2 micron filter. Then aliquots of the solution, containing 0.5 mg/ml peptide in 0.315% m-cresol-1.6% glycerol at pH 8.48, were pipetted into parenteral vials and stoppered.

WO 00/37098

PCT/US99/30395

- (B) A 21.5 ml aliquot of peptide solution in water was mixed with 21.5 ml of 0.63% m-cresol-3.2% glycerol- 0.02 molar L-Lysine pH 8.5 and the final pH was set to 8.48. The solution was passed thru a 0.2 micron filter. Then, aliquots of the solution, containing 0.5 mg/ml peptide in 0.315% m-cresol-1.6% glycerol- 0.01 molar L-Lysine at pH 8.48, were pipetted into parenteral vials and stoppered.
- (C) A 21.5 ml aliquot of peptide solution in water was mixed with 21.5 ml of 0.63% m-cresol-3.2% glycerol- 0.02 molar Tris buffer pH 8.5 and the final pH was set to 8.50. The solution was passed thru a 0.2 micron filter. Aliquots of the solution, containing 0.5 mg/ml peptide in 0.315% m-cresol-1.6% glycerol- 0.01 molar tris at pH 8.50, were pipetted into parenteral vials and stoppered.

One set of vials was held in the refrigerator at 4°C. A second set was held at 30° C (room temperature). A third set was held in the refrigerator, but was taken out three times a day, 5 days a week, and allowed to warm up to room temperature. Then the set was put back in the refrigerator.

At various time points, vials of the three formulations at the three temperature conditions were examined visually, checked with a pH meter, and assayed for main peak purity and total mg/ml of peptide by HPLC. Before assaying on HPLC, the vialled material was spun in centrifuge; the supernatant was injected on the column. The concentration of a freshly prepared solution of Val(8) GLP-1 in 0.01 molar HCl was determined by UV spectroscopy. Standards of known concentration were prepared by diluting this solution; these were injected on HPLC to get a multipoint standard curve. The main peak purity was determined by comparing the main peak area to the total area. The total area (main peak plus related substances) was converted to total mg/ml using the standard curve parameters.

The Table below summarizes a chemical stability study using HPLC to quantify potency, showing that the formulation is stable when there is not a significant loss peptide from solution. %purity is a measure of the stability of the formulation. Chemical degradation of the peptide to soluble related substances results in a lower purity value. The results in this Table show that GLP-1 in TRIS buffer had the highest purity, and hence the highest stability values.

WO 00/37098

PCT/US99/30395

Table 1

Amount of GLP-1	% Purity at 4°	% Purity with starting Temp at 4° with warming to 30°	% Purity at 30°	Buffer added
0.4849	98.13	98.13	98.13	No buffer
0.4795	98.52			No buffer
0.4636		98.26		No buffer
0.4829			95.55	No buffer
0.4801	98.49			No buffer
0.4674		98.43		No buffer
0.4544			94.14	No buffer
0.4327	97.20			No buffer
0.3914		97.82		No buffer
0.3846			87.741	No buffer
0.4567	98.27	98.27	98.27	0.01M lysine
0.4280	98.47			0.01 Lysine
0.4293		98.11		0.01 Lysine
0.4245			94.90	0.01 Lysine
0.4539	98.58			0.01 Lysine
0.4016		98.55		0.01 Lysine
0.4338			93.68	0.01 Lysine
0.3923	96.93			0.01 Lysine
0.3922		96.84		0.01 Lysine
0.4034			85.93	0.01 Lysine
0.4533	98.37	98.37	98.37	0.01 TRIS
0.4602	98.52			0.01 TRIS
0.4340		98.47		0.01 TRIS
0.4374			97.68	0.01 TRIS
0.4709	98.71			0.01 TRIS
0.4531		98.53		0.01 TRIS
0.4377			96.72	0.01 TRIS
0.4569	97.67			0.01 TRIS

WO 00/37098

PCT/US99/30395

Amount of GLP-1	% Purity at 4°	% Purity with starting Temp at 4° with warming to 30°	% Purity at 30°	Buffer added
0.4354		97.81		0.01 TRIS
0.4170			92.13	0.01 TRIS

Example 2-Preparation and Physical Stability Testing

A solution of Val(8) GLP-1 in water was adjusted to pH 8.50 and filtered. The concentration was determined to be 5.28 mg/ml by UV analysis. Aliquots containing 5.00 mg of peptide were pipetted into 10 ml parenteral vials and freeze dried. After freeze drying, the plugs were reconstituted by adding 10.00 ml of either:

- 0.315% m-cresol-1.6% glycerol pH 8.5
- 0.315% m-cresol-1.6% glycerol- 0.005molar Tris buffer pH 8.5
- 0.315% m-cresol-1.6% glycerol- 0.01 molar Tris buffer pH 8.5

Turbidity measurements were measured initially and after 1 month in the refrigerator. All turbidities for the samples (at room temperature) were less than 10 NTU. The initial measurements were made on a Hach 2100N Turbidimeter while the 1 month readings were made on a Hach 2100 AN Turbidimeter.

Buffer	Turbidity (Initial)	Turbidity (1 month)
none	0.4 NTU	4.0 NTU
.005 molar Tris	0.4 NTU	3.6 NTU
.010 molar Tris	0.4 NTU	4.5 NTU

WO 00/37098

PCT/US99/30395

What is claimed is:

1. A shelf-stable pharmaceutical formulation comprising a therapeutically effective amount of a GLP-1 molecule, a pharmaceutically acceptable preservative, and a tonicity modifier, wherein said formulation has a pH that is about 8.2 to about 8.8.

2. The formulation of claim 1, further comprising a buffer.

3. The formulation of claim 2, wherein the buffer is TRIS.

4. The formulation of claim 1, further comprising a surfactant.

5. The formulation of claim 4, wherein the surfactant is Brij-35.

6. The formulation of claim 1, wherein the GLP-1 molecule is an analog of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:

R_1 -X-Glu-Gly¹⁰-Thr-Phe-Thr-Ser-Asp¹⁵-Val-Ser-Ser-Tyr-Leu²⁰-Y-Gly-Gln-Ala-Ala²⁵-
Lys-Z-Phe-Ile-Ala³⁰-Trp-Leu-Val-Lys-Gly³⁵-Arg-R₂ (SEQ ID NO:2)

and a pharmaceutically-acceptable salt thereof, wherein R₁ is His or desamino-histidine, X is Ala, Gly or Val, Y is Glu or Gln, Z is Glu or Gln and R₂ is Gly-OH.

7. The formulation of claim 6, wherein R₁ is L-histidine, X is Val, Y is Glu, Z is Glu, and R₂ is Gly-OH.

8. The formulation of claim 1, wherein the GLP-1 molecule is a derivative of GLP-1 and is selected from the group consisting of a peptide having the amino acid

WO 00/37098

PCT/US99/30395

sequence: $\text{NH}_2\text{-His}^7\text{-Ala-Glu-Gly}^{10}\text{-Thr-Phe-Thr-Ser-Asp}^{15}\text{-Val-Ser-Ser-Tyr-Leu}^{20}\text{-Glu-Gly-Gln-Ala-Ala}^{25}\text{-Lys-Glu-Phe-Ile-Ala}^{30}\text{-Trp-Leu-Val-X}$ (SEQ ID NO:3)

and a pharmaceutically-acceptable salt thereof, wherein X is selected from the group consisting of Lys and Lys-Gly; a pharmaceutically-acceptable lower alkylester of said peptide; and a pharmaceutically-acceptable amide of said peptide selected from the group consisting of amide, lower alkyl amide, and lower dialkyl amide.

9. A method of enhancing the expression of insulin in a mammalian pancreatic β -type islet cell in need of said enhancement, comprising administering to said cell, an effective amount of a shelf-stable pharmaceutical formulation, wherein the formulation comprises a therapeutically effective amount of a GLP-1 molecule, a pharmaceutically acceptable preservative, and a tonicity modifier, and wherein said formulation has a pH that is about 8.2 to about 8.8.

10. The method of claim 9, wherein the formulation further comprises a buffer.

11. The method of claim 10, wherein the buffer is TRIS.

12. The method of claim 9, wherein the formulation further comprises a surfactant.

13. The method of claim 12, wherein the surfactant is Brij-35.

14. The method of claim 9, wherein the GLP-1 molecule is an analog of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:

$\text{R}_1\text{-X-Glu-Gly}^{10}\text{-Thr-Phe-Thr-Ser-Asp}^{15}\text{-Val-Ser-Ser-Tyr-Leu}^{20}\text{-Y-Gly-Gln-Ala-Ala}^{25}\text{-Lys-Z-Phe-Ile-Ala}^{30}\text{-Trp-Leu-Val-Lys-Gly}^{35}\text{-Arg-R}_2$ (SEQ ID NO:2)

WO 00/37098

PCT/US99/30395

and a pharmaceutically-acceptable salt thereof, wherein R_1 is His or desamino-histidine, X is Ala, Gly or Val, Y is Glu or Gln, Z is Glu or Gln and R_2 is Gly-OH.

15. The method of claim 14, wherein R_1 is L-histidine, X is Val, Y is Glu, Z is Glu, and R_2 is Gly-OH.

16. The method of claim 9, wherein the GLP-1 molecule is a derivative of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:

$\text{NH}_2\text{-His}^7\text{-Ala-Glu-Gly}^{10}\text{-Thr-Phe-Thr-Ser-Asp}^{15}\text{-Val-Ser-Ser-Tyr-Leu}^{20}\text{-Glu-Gly-Gln-Ala-Ala}^{25}\text{-Lys-Glu-Phe-Ile-Ala}^{30}\text{-Trp-Leu-Val-X}$ (SEQ ID NO:3)

and a pharmaceutically-acceptable salt thereof, wherein X is selected from the group consisting of Lys and Lys-Gly; a pharmaceutically-acceptable lower alkylester of said peptide; and a pharmaceutically-acceptable amide of said peptide selected from the group consisting of amide, lower alkyl amide, and lower dialkyl amide.

17. A method for treating diabetes comprising administering to a patient in need of such treatment an effective amount of a shelf-stable pharmaceutical formulation, wherein the formulation comprises a therapeutically effective amount of a GLP-1 molecule, a pharmaceutically acceptable preservative, and a tonicity modifier, and wherein said formulation has a pH that is about 8.2 to about 8.8.

18. The method of claim 17, wherein the formulation further comprises a buffer.

19. The method of claim 18, wherein the buffer is TRIS.

20. The method of claim 17, wherein the formulation further comprises a surfactant.

WO 00/37098

PCT/US99/30395

21. The method of claim 20, wherein the surfactant is Brij-35.

22. The method of claim 17, wherein the GLP-1 molecule is an analog of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence:

R_1 -X-Glu-Gly¹⁰-Thr-Phe-Thr-Ser-Asp¹⁵-Val-Ser-Ser-Tyr-Leu²⁰-Y-Gly-Gln-Ala-Ala²⁵-
Lys-Z-Phe-Ile-Ala³⁰-Trp-Leu-Val-Lys-Gly³⁵-Arg-R₂ (SEQ ID NO:2)

and pharmaceutically-acceptable salts thereof, wherein R₁ is His or desamino-histidine, X is Ala, Gly or Val, Y is Glu or Gln, Z is Glu or Gln and R₂ is Gly-OH.

23. The method of claim 22, R₁ is L-histidine, X is Val, Y is Glu, Z is Glu, and R₂ is Gly-OH.

24. The method of claim 17, wherein the GLP-1 molecule is a derivative of GLP-1 and is selected from the group consisting of a peptide having the amino acid sequence: NH₂-His⁷-Ala-Glu-Gly¹⁰-Thr-Phe-Thr-Ser-Asp¹⁵-Val-Ser-Ser-Tyr-Leu²⁰-Glu-Gly-Gln-Ala-Ala²⁵-Lys-Glu-Phe-Ile-Ala³⁰-Trp-Leu-Val-X (SEQ ID NO:3)

and pharmaceutically-acceptable salts thereof, wherein X is selected from the group consisting of Lys and Lys-Gly; a pharmaceutically-acceptable lower alkylester of said peptide; and a pharmaceutically-acceptable amide of said peptide selected from the group consisting of amide, lower alkyl amide, and lower dialkyl amide.

25. The formulation of claim 1, further comprising a long-acting insulin agent.

26. A method of providing meal-time glycemic control and basal glycemic control with a single injection comprising administering to a patient in need thereof an effective amount of a shelf-stable pharmaceutical formulation, wherein the formulation

WO 00/37098

PCT/US99/30395

comprises a therapeutically effective amount of a GLP-1 molecule, a long acting insulin agent, a pharmaceutically acceptable preservative, and a tonicity modifier, wherein said formulation has a pH that is about 8.2 to about 8.8.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 38/26, 38/28, 47/10, 47/26, A61P 3/10 // (A61K 38/28, 38:26)	A1	(11) International Publication Number: WO 00/37098 (43) International Publication Date: 29 June 2000 (29.06.00)
(21) International Application Number: PCT/US99/30395 (22) International Filing Date: 21 December 1999 (21.12.99) (30) Priority Data: 60/113,499 22 December 1998 (22.12.98) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BRADER, Mark, L. [NZ/US]; 6465 North Park Avenue, Indianapolis, IN 46220 (US). PEKAR, Allen, H. [US/US]; 5354 North Park Avenue, Indianapolis, IN 46220 (US). (74) Agent: MACIAK, Ronald, S.; Eli Lilly and Company, Lilly Corporate Center, Drop Code1501, Indianapolis, IN 46285 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: SHELF-STABLE FORMULATION OF GLUCAGON-LIKE PEPTIDE-1 (57) Abstract Glucagon-like peptide-1 (GLP-1) has been shown to be useful in the treatment of diabetes. The invention encompasses a shelf stable formulation that comprises a therapeutically effective amount of GLP-1, a pharmaceutically acceptable preservative, and a tonicity modifier, and that has a pH between about 8.2 to about 8.8.		

Please type a plus sign (+) inside this box



09868974 062001

PTO/SB/01 (8-96) (MODIFIED)

Approved for use through 9/30/98. OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION

☒ Declaration Submitted with Initial Filing
☐ Declaration Submitted after Initial Filing

Attorney Docket Number	X-11869
First Named Inventor	Mark Laurence Brader, et al.
COMPLETE IF KNOWN	
Application Number	
Filing Date	
Group Art Unit	
Examiner Name	

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

SHELF-STABLE FORMULATION OF GLUCAGON-LIKE PEPTIDE-1

the specification of which

☐ is attached hereto

OR

☒ was filed on (MM/DD/YYYY) 12/21/1999 as United States Application Number or PCT International

Application Number PCT/US99/30395 and was amended on (MM/DD/YYYY) (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37 Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached	
				YES	NO
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority sheet attached hereto:

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional applications(s) listed below.

Application Number(s) 60/113,499	Filing Date (MM/DD/YYYY) 12/22/1998	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority sheet attached hereto.
-------------------------------------	--	--

Please type a plus sign (+) inside this box



PTO/SB/01 (8-96) (MODIFIED)

Approved for use through 9/30/98. OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

DECLARATION

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s), or § 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority sheet attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Attorney Name	Reg. No.
Arvie J. Anderson	45,263
Lynn D. Apelgren	45,341
Robert A. Armitage	27,417
Brian P. Barrett	39,597
Michael T. Bates	34,121
Roger S. Benjamin	27,025
William R. Boudreaux	35,796
Steven P. Caltrider	36,467
Paul R. Cantrell	36,470
Charles E. Cohen	34,565
Robert A. Conrad	32,089
Donald L. Corneglio	30,741
Gregory A. Cox	P47,504
Paul R. Darkes	33,862
Paula K. Davis	P47,517
Elizabeth A. Dawalt	44,646
John C. Demeter	30,167
Manisha A. Desai	43,585
Joanne Longo Feeney	35,134
Paul J. Gaylo	36,808
Francis O. Ginah	44,712
Amy E. Hamilton	33,894
Frederick D. Hunter	26,915
Thomas E. Jackson	33,064
Charles T. Joyner	30,466
Gerald P. Keleher	43,707

Attorney Name	Reg. No.
James J. Kelley	41,888
Paul J. Koivuniemi	31,533
Robert E. Lee	27,919
Kirby Lee	P47,744
James P. Leeds	35,241
Nelsen L. Lentz	38,537
Janet T. McClain	36,863
Douglas K. Norman	33,267
Arleen Palmberg	40,422
Raymond S. Parker, III	34,893
Thomas G. Plant	35,784
Edward Prein	37,212
James J. Sales	33,773
Michael J. Sayles	32,295
Robert L. Sharp	45,609
David M. Stemerick	40,187
Mark J. Stewart	43,936
Robert D. Titus	40,206
Robert C. Tucker	45,165
Tina M. Tucker	47,145
MaCharri Vorndran-Jones	36,711
Gilbert T. Voy	43,972
Thomas D. Webster	39,872
Lawrence T. Welch	29,487
Alexander Wilson	P45,782

☐ Additional registered practitioner(s) named on a supplemental sheet attached hereto.

Direct all correspondence to:

Name	ELI LILLY AND COMPANY				
Address	ATTN: Mark J. Stewart				
Address	LILLY CORPORATE CENTER/DC1104				
City	INDIANAPOLIS	State	INDIANA	ZIP	46285
Country		Telephone	(317) 276-0280	Fax	(317) 276-3861

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor:

☐ A Petition has been filed for this unsigned inventor

Given Name	Mark	Middle Name	Laurence	Family Name	Brader	Suffix e.g. Jr.	
Inventor's Signature	Mark Laurence Brader				Date	6/20/01	
Residence: City	Indianapolis	IN	State	IN	Country	US	Citizenship
Address	5858 Forest Lane						
Post Office Address	SAME AS ABOVE						
City	Indianapolis	State	IN	Zip	46220	Country	US

☒ Additional Inventors are being named on supplement sheet(s) attached hereto.

09268974 062003

Please type a plus sign (+) inside this box ☐

PTO/SB/01 (8-96) (MODIFIED)

Approved for use through 9/30/98. OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

DECLARATION

Name of Additional Joint Inventor, if any:

☐ A Petition has been filed for this unsigned inventor

Given Name	Allen	Middle Name	Howard	Family Name	Pekar	Suffix e.g. Jr.	
Inventor's Signature	<i>Allen Howard Pekar</i>					Date	6-19-2001
Residence: City	Indianapolis	IN	State	IN	Country	US	Citizenship
Address	5354 North Park Avenue						
Post Office Address	SAME AS ABOVE						
City	Indianapolis	State	IN	Zip	46220	Country	US

09/868974

X-11869.ST25.txt
SEQUENCE LISTING

<110> Brader, Mark L.
Pekar, Allen H.

<120> SHELF-STABLE FORMULATION OF GLUCAGON-LIKE PEPTIDE-1

<130> X-11869

<140> US 09/868,974

<141> 1999-12-21

<150> PCT/US99/30395

<151> 1999-12-21

<150> US 60/113,499

<151> 1998-12-22

<160> 3

<170> PatentIn version 3.1

<210> 1

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<220>

<221> MISC_FEATURE

<222> (2)..(2)

<223> Xaa at position 2 is Ala, Gly or Val

X-11869.ST25.txt

<220>

<221> MISC_FEATURE

<222> (15)..(15)

<223> xaa at position 15 is Glu or Gln

<220>

<221> MISC_FEATURE

<222> (21)..(21)

<223> xaa at position at 21 is Glu or Gln

<400> 1

His Xaa Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Xaa Gly
 1 5 10 15

Gln Ala Ala Lys Xaa Phe Ile Ala Trp Leu Val Lys Gly Arg Gly
 20 25 30

<210> 2

<211> 29

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<220>

<221> MISC_FEATURE

<222> (29)..(29)

<223> xaa at position 29 is Gly or is absent

<400> 2

His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Xaa
 20 25

<210> 3

X-11869.ST25.txt

<211> 30

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<220>

<221> MISC_FEATURE

<222> (19)..(19)

<223> Xaa at position 19 is Lys or Arg

<220>

<221> MISC_FEATURE

<222> (30)..(30)

<223> Xaa at position 30 is Gly or is absent

<400> 3

Ala	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Val	Ser	Ser	Tyr	Leu	Glu	Gly	Gln
1				5					10					15	

Ala	Ala	Xaa	Glu	Phe	Ile	Ala	Trp	Glu	Val	Lys	Gly	Arg	Xaa
			20					25					30